

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

5,000

Open access books available

125,000

International authors and editors

140M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Procoagulant Platelets

Andaleb Kholmukhamedov

Abstract

There are two well-known subpopulations of activated platelets: pro-aggregatory and procoagulant. Procoagulant platelets represent a subpopulation of activated platelets, which are morphologically and functionally distinct from pro-aggregatory ones. Although various names have been used to describe these platelets in the literature (CoaT, CoaTed, highly activated, ballooned, capped, etc.), there is a consensus on their phenotypic features including exposure of high levels of phosphatidylserine (PSer) on the surface; decreased aggregatory and adhesive properties; support of active tenase and prothrombinase complexes; maximal generation by co-stimulation of glycoprotein VI (GPVI) and protease-activated receptors (PAR). In this chapter, morphologic and functional features of procoagulant platelets, as well as the mechanisms of their formation, will be discussed.

Keywords: procoagulant, platelet, mitochondria, necrosis, apoptosis, coagulation

1. Introduction

Blood has different components, like plasma, red blood cells (RBC), white blood cells (WBC), and platelets. Platelets, although being only “tiny fragments of megakaryocytes (mother cell),” are essential for life. We need them, together with about two dozen of coagulation factors, to keep all that nutrient-rich liquid plasma, infection-fighting WBCs, and oxygen-carrying RBCs in our bodies in the case of trauma and bleeding. Platelets and coagulation get activated immediately upon being exposed to things they normally do not have contact with (e.g., during blood vessel wall rupture, collagen gets exposed and activates platelets). Upon activation, platelets form a mesh-like structure using another plasma protein called fibrinogen as bridges between them. This process forms a so-called “unstable clot.” At later stages of blood clotting, generated thrombin converts soluble fibrinogen to insoluble fibrin, stabilizing the initial platelet plug. These processes are collectively called hemostasis. Every aspect of hemostasis has its history. Although exploring history can sometimes be tedious, studying the bridge between past and present is essential in basic understanding of the subject, the subject of procoagulant platelets in this case.

Unlike WBCs, which exist as functionally and morphologically distinct subpopulations, it has been thought for years that platelets are rather simple in their function, being just “cell particles.” It was later revealed that platelets, although lacking nuclear material, are indeed very complex not only in their nature but also in function. Nowadays, the existence of two different subpopulations of activated platelets, pro-aggregatory and procoagulant, is a widely accepted fact. Pro-aggregatory platelets, historically known as activated, have been a major focus since the initiation of platelet research. The history of procoagulant platelets, on the other

hand, can be subdivided into two periods. The first one is the discovery of platelet procoagulant functionality, whereas in the second period, procoagulant platelets were discovered and characterized as a distinct subpopulation of activated platelets.

In 1912, long before we learned that prothrombinase supporting platelets are distinct from pro-aggregatory ones, Howell discovered that unsaturated cephalin is a phospholipid factor that triggers clotting [1]. This discovery was followed by decades of controversial results of what phospholipid or what mixture of phospholipids is responsible for this effect until, in 1960, Karl Slotta demonstrated the presence of phosphatidylserine (PSer) is an absolute requirement [2]. This discovery established the central role of PSer in prothrombinase activity, which in a way, paused research activities in the procoagulant platelet field.

The history was resumed when two decades later, Bevers and colleagues spectrophotometrically showed that thrombin and collagen co-stimulated platelets possessing about 5-fold higher prothrombinase activity than those stimulated with thrombin or collagen alone. This effect was even more pronounced in the presence of exogenous factor Va [3]. Later that year, the same group confirmed their spectrophotometric observations in one-stage prothrombinase assay [4]. In 1985, Rosing et al. more thoroughly described the role of platelet PSer exposure in prothrombin (FII) and factor X (FX) activation [5]. They revealed that collagen and thrombin co-stimulation significantly increases prothrombinase and tenase activity, while this increase was abolished in the presence of phospholipase A₂ (PLA₂) from *N. naja*. Also observing no cellular lysis in these conditions, they concluded that PSer exposure, which is followed by thrombin and collagen co-stimulation, is responsible for this phenomenon. In 1993, with the discoveries of annexin V and advancement in flow cytometry, Dachary-Prigent et al. established a protocol to detect PSer exposing platelets, the methodology that has overtime become fundamental in procoagulant platelet field and has been widely used since then [6]. In 1997 Heemskerk and colleagues discovered that a percentage of platelets adhering to collagen, but not fibrinogen, balloon and expose PSer [7]. In parallel to these findings, there was a series of publications determining the extent and ultrastructure of PSer exposure, as well as the essential role of calcium in this process [8–10].

Although numerous studies described platelet procoagulant function in response to dual stimulation [3, 5, 11, 12], a breakthrough discovery that identified procoagulant platelets as a distinct subpopulation of activated platelets is the work by Alberio et al., where they demonstrated that only a certain percentage of activated platelets retain factor Va (FVa) on their surface [13]. This subpopulation was generated by co-stimulation with convulxin/collagen (GPVI agonists) and thrombin; hence they were named CoaT platelets. A few years down the road, Kulkarni and Jackson introduced a new term—‘sustained calcium-induced platelet morphology (SCIP)’ by discovering the fact that procoagulant platelets require prolonged elevation in cytosolic calcium to form [14]. Diversifying terminology of procoagulant platelets did not stop there, in 2005 revealing that procoagulant platelets retain multiple α -granule proteins on their surface in transglutaminase-dependent manner led to the introduction of a new term—‘coated’ (by α -granule proteins) platelets [15, 16]. Further studies by Panteleev et al. revealed that this subpopulation of activated platelets binds high levels of factors IXa and Xa [17]. In 2008, Jobe et al., characterizing molecular mechanisms of procoagulant platelet formation, introduced another term—highly activated platelets [18]. There are few more terms used in the literature like ballooned [7], ballooned and procoagulant-spread [19], and super-activated platelets [20].

Just like in a famous Indian parable, where blind men try to describe an elephant they have never encountered before by touching different parts of it, researchers have been describing different (morphological and functional) features of

procoagulant platelets and introducing different terms based on their discoveries. Whereas indeed, everyone has been describing “different parts of the same elephant.”

2. Mechanisms of procoagulant platelet formation

The discovery of procoagulant platelets as a distinct subpopulation of activated platelets at the beginning of the twenty-first century triggered research activities into cellular and molecular mechanisms of their formation. As shown by Alberio et al. and confirmed in later studies, procoagulant platelets are maximally generated upon co-stimulation of glycoprotein VI and PAR1/4. In 2005, Jobe and colleagues discovered that the absence of FcR γ , a key component responsible for glycoprotein VI signaling, ablates procoagulant platelet formation almost to baseline levels, evidencing GPVI stimulation is the major component of their generation [21]. The same year, Remenyi et al. demonstrated the role of mitochondrial permeability transition (MPT) in procoagulant platelet formation [16]. MPT is a Ca²⁺-dependent molecular process that leads to mitochondrial swelling and cell death [22, 23]. During the onset of MPT, large pores are formed on the mitochondrial inner membrane making it non-specifically permeable to all solutes and molecules of molecular weight up to 1500 Da [24, 25]. It is a very well-known fact that mitochondrial Ca²⁺ overload can induce MPT, although the structure of MPT pore remains unknown.

Ca²⁺, being a key signaling molecule in most cells, is important for many processes including platelet shape change and integrin α 2b β 3 activation [26, 27]. In resting platelets, free Ca²⁺ is tightly regulated and maintained at about 100 nM in both the cytosol and mitochondria through the action of Ca²⁺-ATPases in the plasma and cell membranes. Thus, platelet cytosolic-free Ca²⁺ is substantially lower than the blood Ca²⁺ levels, which are around 2 mM. With stimulation, however, cytosolic Ca²⁺ increases instantaneously. As outlined in **Figure 1**, this increase is mediated by activated phospholipase C (PLC). There are two major isoforms of PLC in human platelets, PLC β and PLC γ . PLC β is only activated downstream of Gq protein-coupled receptors (GPCRs). Whereas, PLC γ is activated downstream of numerous receptors like GPVI, glycoprotein Ib-IX complex (GPIb-IX), Fc γ receptor IIa (Fc γ RIIa), and C-type lectin-like receptor 2 (CLEC-2). As illustrated in **Figure 1**, both isoforms of PLC induce the release of Ca²⁺ from the dense tubular system (DTS) as well as activating transient receptor potential channel 3 (TRPC3). DTS release of Ca²⁺, in turn, triggers its extracellular entry through store-operated calcium entry (SOCE). For some time, it remained mysterious on how the release of DTS calcium stores into the cytosol induces more Ca²⁺ to flow into the cell, further increasing its cytosolic concentrations. However, with the discovery of core components of SOCE, everything falls into place. STIM1 and ORAI1 are parts of the same complex. When STIM1 senses drop in Ca²⁺ levels within the DTS, it signals to ORAI1 located on a cell membrane to allow Ca²⁺ passage from extracellular space into the cell. As evidenced by the study in the early 2000s, increased cytosolic Ca²⁺ levels are one of the requirements of procoagulant formation [14]. Mitochondria, being the major and perhaps the only Ca²⁺ buffering system within platelets, equilibrate increased cytosolic Ca²⁺ following PAR and GPVI co-stimulation. The ensuing fate of mitochondrial Ca²⁺ overload is the opening of the mitochondrial permeability transition pore (MPTP), followed by PSer exposure and integrin deactivation (**Figure 1**), ultimately leading to a cell death. However, in this case cell death shall be considered physiologic considering the essential role of procoagulant platelets in hemostasis.




Figure 1.

Molecular mechanisms of physiologic agonist-induced procoagulant platelet formation. In resting platelets, low cytosolic Ca^{2+} is maintained by cell and plasma membrane Ca^{2+} -ATP-ases. Co-stimulation of protease-activated receptor 1/4 and glycoprotein VI leads to a sustained increase of cytosolic Ca^{2+} through activation of $\text{PLC}\beta$ and $\text{PLC}\gamma$, respectively. Increased cytosolic Ca^{2+} drives mitochondrial Ca^{2+} entry through the mitochondrial calcium uniporter complex. The resulting increase in mitochondrial Ca^{2+} opens mitochondrial permeability transition pore, which in turn leads to necrotic cell death and exposure of phosphatidylserine accompanied by integrin inactivation. DTS, dense tubular system; GPVI, glycoprotein VI; MCU, mitochondrial calcium uniporter; MPTP, mitochondrial permeability transition pore; PAR, protease-activated receptor; PSer, phosphatidylserine; SOCE, store-operated calcium entry; TRPC3, transient receptor potential channel 3. Adapted from [61] with modifications.

It brings us to another aspect of procoagulant platelet research, debated since their discovery, which is whether procoagulant platelets are necrotic or apoptotic. Cell death, both necrosis and apoptosis, is an essential event in the normal life of many cells in the human body, including platelets. Apoptosis, or programmed cell death, occurs in many organs throughout human lifetime. Necrosis, although considered to be mostly a catastrophic uncontrolled cell death, can also occur physiologically, as in the shedding of decidual endometrium during human menses. Necrosis and apoptosis differ in many aspects. The ultimate event during necrotic cell death is the osmotic swelling of the cell followed by the rupture of a cell membrane, whereas in apoptosis, cell shrinkage with preserved cell membrane is evident in later stages of this process.

Although the proposal that PSer is not homogeneously distributed on the surface of activated platelets was made back in 1985 [5], it took three decades to visualize that experimentally due to the complex nature of procoagulant platelets. In 2016, Podoplelova et al. presented a detailed structure of procoagulant platelets and cellular changes that give them the characteristic morphology. They elegantly demonstrated how the balloon is blown out from the platelet, leaving it as a “cap” [28]. The existence of bulges in a phospholipid bilayer, known as the open canalicular system (OCS), is essential, as it provides platelet a fair amount of surface reserve for ballooning. Morphologic appearance, in this case, resembles the classical osmotic swelling of necrotic cell death, as schematically presented on the left side of **Figure 2**. Another feature of necrosis, which is a collapse of energy production, is present in procoagulant platelets, as they lack energy-requiring contractile function [18]. Moreover, depletion of oxidative




Figure 2.
Cellular mechanisms of platelet cell death. Cyt C, cytochrome C; GPVI, glycoprotein VI; mPTP, mitochondrial permeability transition pore; PSer, phosphatidylserine. PSer exposure is shown in red.

phosphorylation due to MPT, deactivates flippases that normally maintain the asymmetry of phospholipids. After the onset of MPT, contents of mitochondrial matrix get released to the cytosol leading to the second wave of cytosolic Ca^{2+} increase. This in turn leads to scramblase activation. Synergistically with deactivated flippases, scramblase performs its raison d'être, which is flipping negatively charged PSer out in order to equilibrate its concentration. And finally, recent work demonstrated that platelets undergoing cell death through necrotic pattern and not apoptotic are functionally procoagulant as measured by prothrombinase support [29].

Platelet apoptosis (**Figure 2**, right side), on the other hand, has been widely implicated in platelet lifespan via the action of the intrinsic mitochondrial apoptosis pathway [30, 31]. Normally, antiapoptotic members of the BCL-2 family (e.g., BCL-xL) restrain the activity of proapoptotic Bax and Bak proteins, which are present within the cytosolic fraction of a platelet. When BCL-xL wears off, oligomerized Bax and Bak translocate to the mitochondrial outer membrane and permeabilize it, which leads to cytochrome C leakage into the cytosol, triggering apoptosome formation and eventually cell death. This is supported by the fact that the genetic ablation of murine BCL-xL reduces platelet life span from about 5 days to 5 hours [32, 33].

Thus, procoagulant platelets are indeed necrotic, while apoptosis is essential in platelet clearance. It should, however, be mentioned that the presence of PSer on the surface of a platelet is a signal for the reticuloendothelial system in the spleen and liver for clearance. Therefore, any procoagulant platelet that happened to escape the site of the active hemostatic process will be cleared out of the system by liver or spleen, just like aged apoptotic platelets do.

3. Functions of procoagulant platelets

The physiological relevance of procoagulant platelets had been questioned for a long time. In recent years, however, after the demonstration of a procoagulant platelet being predictive of bleeding or ischemic complications in patients with coronary artery disease, brain hemorrhage, traumatic brain injury, stroke, etc. [34–41] it is gaining more and more recognition. The importance of this subpopulation is further highlighted on a novel *ex vivo* model (which integrates all the core components of hemostasis [42]), where pharmacologic inhibition of platelet transition to a procoagulant state without affecting pro-aggregatory phenotype results in a decreased thrombus stability [43].

It was initially thought that the only function of procoagulant platelets is to support coagulation. However, with recent advances in the field, we learn that depending on their localizations, procoagulant platelets can play different functions within the thrombus. For general consideration, these two functionalities will be discussed separately here.

3.1 Coagulation support

Coagulation, a cascade of serine protease enzymatic reactions, is achieved by cleaving fibrinogen to fibrin, which transforms blood from a liquid to a gel-like state. Although platelet contribution to coagulation has been known for decades, the exact role of platelet phospholipids has been a matter of major debate. Dependence of hemostasis on biological membranes is very extensive, ranging from subendothelial membranes triggering coagulation and platelet activation to procoagulant platelet surface assembling tenase and prothrombinase complexes. It is very well known that at least two coagulation reactions are highly dependent on phospholipid surface, namely the activation of factor X and prothrombin by intrinsic tenase and prothrombinase, respectively. Both intrinsic tenase and prothrombinase are composed of the serine protease (FIXa for tenase and FXa for prothrombinase) and its protein cofactor (FVIIIa for tenase and FVa for prothrombinase). It is important to know that although both proteases alone are capable of activating their substrates, the presence of cofactors profoundly amplifies the catalytic process for up to 10,000 fold.

The current understanding of this process, based on numerous studies including mathematical modeling, is that phospholipid surface increases the rate of reactions by increasing the local concentration of coagulation factors, and thus increasing the probability of their interaction [28, 44–50]. This increase in a local concentration of factors, essential for tenase and prothrombinase complexes, is accomplished by the interaction of negatively charged gamma-carboxyglutamic acid (GLA) residues of the coagulation factors with negatively charged phosphatidylserine on the surface of procoagulant platelets. It is supported by the fact that GLA domain-containing (also known as vitamin K-dependent) factors predominantly bind to a procoagulant subpopulation of activated platelets [17, 28, 29]. Both enzymatic factors for intrinsic tenase (FIX) and prothrombinase (FX) are GLA domain-containing proteins and bind to PSer in a calcium-dependent manner. Whereas cofactors (FVIII and FV) structurally are not GLA proteins and bind to a procoagulant surface by different mechanisms.

In the case of FVIII, it has been shown by Gilbert et al. that it is not specific to a procoagulant surface [51]. It was further confirmed by Podoplelova et al. in their efforts exploring procoagulant platelet characteristics, they demonstrated that both (pro-aggregatory and procoagulant) subpopulations of activated platelets bind FVIIIa [28]. The fact that PS increases the catalytic activity of the intrinsic tenase

complex by about 1500-fold [52] can be explained by FIXa's specificity to PSer. As outlined in **Figure 3A**, initially FIXa and FX bind PSer. This binding reaction is calcium-dependent for all GLA domain-containing proteins. FIXa possesses enzymatic activity to convert X to Xa, whereas in the presence of its cofactor (FVIIIa), the catalytic activity raises 100,000-fold.

As for the FV—there are two different probabilities of binding to the procoagulant surface, as demonstrated in **Figure 3B**. In the first one, prothrombinase forms by the surface Xa reacting with factor Va in solution. Whereas, in the second case, both FVa and FXa form binary complexes with PSer first, and then lipid-protein rearrangement leads to prothrombinase formation [49]. The addition of exogenous FVa to procoagulant platelets increases the velocity of prothrombinase reaction [4], whereas in the absence of exogenous FVa, procoagulant platelets are still prothrombinase active [53]. These findings indicate that both pathways are physiologically important.

3.2 Limiting thrombus growth

During hemostasis, there is a time point when a thrombus needs to stop growing in order not to occlude the lumen of the vessel, which may compromise the blood supply to an end organ. Stoppage of the thrombus growth is probably the most intriguing part of thrombogenesis, although being the most understudied at the same time. When coagulation is initiated by subendothelial tissue factor, it leads to the formation of small amounts of thrombin. Thrombin then, via a positive feedback mechanism, by activating FXI and FVIII triggers the contact activation pathway and amplifies its production. This self-accelerating process is essential within the hemostatic plaque as it is the only way to overcome the anticoagulant nature of plasma, which is due to the presence of antithrombin III (ATIII). Besides the presence of ATIII in the active form, other mechanisms limit coagulation beyond its border zone. Another one of high importance is the presence of thrombomodulin, an integral membrane protein that is expressed ubiquitously on the surface of endothelial cells. Thrombomodulin converts procoagulant thrombin to an anticoagulant enzyme. Not only these mechanisms limit coagulation to the injured site but also degrade activated factors that happened to escape the hemostatic plaque.




Figure 3. Factor X activation (A) and prothrombinase assembly (B) on the “cap” of procoagulant platelet. (A). Initially FIXa and FX are bound to PSer on the surface of procoagulant platelet in calcium-dependent manner (①), whereas co-factor VIII binding is not specific to PSer and does not require calcium (②). FIXa possesses enzymatic activity to convert X to Xa (③). Although reaction ③ is very slow in the absence of factor FVIIIa, it is efficient to generate small amounts of FXa. Initial FXa then activates VIII to FVIIIa (④), this leads to intrinsic tenase complex assembly, which in turn amplifies Xa formation (⑤). (B). There are two possibilities for the assembly of prothrombinase on the surface of procoagulant platelet. Pathway A forms prothrombinase by the surface Xa reacting with factor Va in solution. Pathway B depicts the possibility for prothrombinase complex formation by already bound Xa and Va. In this case, both FVa and FXa form binary complexes with PSer first and then lipid-protein rearrangement leads to prothrombinase formation.

These processes outline the general principles of limiting coagulation to its border zone. However, not only coagulation but also processes of platelet adhesion, aggregation, and activation have to stop in order at a certain timepoint. How and when thrombus stops growing concerning cellular component had been a mystery for a long time.

In 2007, Maroney et al. demonstrated that procoagulant platelets express active tissue factor pathway inhibitor (TFPI) on their surface [54]. But why platelets expressing procoagulant PSer would also need a strong anticoagulant (TFPI) on




Figure 4. Schematic representation of platelet translocation within the evolving hemostatic plug. After vascular injury, platelets adhere and aggregate at the wound site. A subpopulation of platelets within the hemostatic plug transitions to a procoagulant state. Procoagulant and pro-aggregatory platelets have opposing vectors of translocation. Clot retraction, driven by platelets with a pro-aggregatory phenotype, squeezes procoagulant platelets to the periphery. Adapted from [62] with minor modification.

their surface remained a question for about a decade until in 2019 a breakthrough work by Nechipurenko et al. characterized a phenomenon, together with its mechanisms, of procoagulant platelets translocating to the thrombus periphery [55]. They demonstrated that during clot retraction procoagulant platelets are squeezed out to the periphery of the thrombus, as shown in **Figure 4**. Another important study in this context is the 2008 work by Jobe et al., which demonstrated that procoagulant platelets do not possess contractile function [18]. This explains why pro-aggregatory and procoagulant platelets have different translocation vectors within the hemostatic plaque (**Figure 4**). Being bound to the thrombus by its “cap,” procoagulant platelets do not get detached but rather are squeezed out during pro-aggregatory platelets contraction. At the luminal surface of the hemostatic plaque, procoagulant platelets limit its further growth not only by expressing low adhesive and aggregatory surfaces [56–58], but also with TFPI terminating any extrinsic tenase and prothrombinase activity.

The fact of procoagulant platelet being non-adhesive, however, gives rise to a legitimate question. If procoagulant platelets are not capable of adhesion and aggregation, how do they get attracted to a thrombus? It turns out to be, as described in recent studies, that procoagulant platelets do not form *de novo* from a resting state, but rather temporally transitioning from pro-aggregatory phenotype by the onset of the MPT and regulated necrosis within a hemostatic plaque [59, 60].

Many are aware of an ancient Roman god of duality — Janus, who had two faces as he was able to look to the future and the past. Procoagulant platelets can also demonstrate a functional duality by supporting coagulation on one side and limiting thrombus growth on the other, making them be “Janus” of hemostasis.

Conflict of interest

The author has no conflict of interest to disclose.

Abbreviations

APC	activated protein C
ATIII	antithrombin III
Cyt C	cytochrome c
DTS	dense tubular system
FcRγ	FC receptor gamma
FV	factor V
FVa	activated factor V
FVIII	factor VIII
FVIIIa	activated factor VIII
FIX	factor IX
FIXa	activated factor IX
FX	factor X
FXa	activated factor X
GPIb-IX	glycoprotein Ib and IX complex
GPVI	glycoprotein VI
MCU	mitochondrial calcium uniporter
MPT	mitochondrial permeability transition
MPTP	mitochondrial permeability transition pore
OCS	open canalicular system
PAR	protease-activated receptor

PC	protein C
PLA ₂	phospholipase A ₂
PS	protein S
PSer	phosphatidylserine
RBC	red blood cell
SOCE	store-operated calcium entry
TF	tissue factor
TFPI	tissue factor pathway inhibitor
TRPC3	transient receptor potential channel 3
WBC	white blood cell

IntechOpen

Author details

Andaleb Kholmukhamedov
Versiti Blood Research Institute, Milwaukee, Wisconsin, USA

*Address all correspondence to: akholmukhame@ilww.com

IntechOpen

© 2020 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. 

References

- [1] Howell W. Nature and action of thromboplastic (zymoplastic) substances of tissues. *American Journal of Physiology*. 1912;**31**:1-21
- [2] Slotta K. Thromboplastin I. Phospholipid moiety of Thromboplastin. *Proceedings of the Society for Experimental Biology and Medicine*. January 1960;**103**:53-56
- [3] Bevers EM, Comfurius P, van Rijn JL, Hemker HC, Zwaal RF. Generation of prothrombin-converting activity and the exposure of phosphatidylserine at the outer surface of platelets. *European Journal of Biochemistry*. 1982;**122**(2):429-436
- [4] Bevers EM, Comfurius P, Hemker HC, Zwaal RFA. On the clot-promoting activity of human platelets in a one-stage prothrombinase assay. *Pathophysiology of Haemostasis and Thrombosis*. 1982;**12**(3):268-274
- [5] Rosing J, van Rijn JL, Bevers EM, van Dieijen G, Comfurius P, Zwaal RF. The role of activated human platelets in prothrombin and factor X activation. *Blood*. 1985;**65**(2):319-332
- [6] Dachary-Prigent J, Freyssinet JM, Pasquet JM, Carron JC, Nurden AT. Annexin V as a probe of aminophospholipid exposure and platelet membrane vesiculation: A flow cytometry study showing a role for free sulfhydryl groups. *Blood*. 1993;**81**(10):2554-2565
- [7] Heemskerk JW, Vuist WM, Feijge MA, Reutelingsperger CP, Lindhout T. Collagen but not fibrinogen surfaces induce bleb formation, exposure of phosphatidylserine, and procoagulant activity of adherent platelets: Evidence for regulation by protein tyrosine kinase-dependent Ca²⁺ responses. *Blood*. 1997;**90**(7):2615-2625
- [8] Trotter PJ, Orchard MA, Walker JH. Thrombin stimulates the intracellular relocation of annexin V in human platelets. *Biochimica et Biophysica Acta*. 1994;**1222**(2):135-140
- [9] Trotter PJ, Orchard MA, Walker JH. Ca²⁺ concentration during binding determines the manner in which annexin V binds to membranes. *The Biochemical Journal*. 1995;**308** (Pt 2):591-598
- [10] Tzima E, Poujol C, Nurden P, Nurden AT, Orchard MA, Walker JH. Annexin V relocates to the periphery of activated platelets following thrombin activation: An ultrastructural immunohistochemical approach. *Cell Biology International*. 1999;**23**(9):629-635
- [11] Smeets EF, Heemskerk JW, Comfurius P, Bevers EM, Zwaal RF. Thapsigargin amplifies the platelet procoagulant response caused by thrombin. *Thrombosis and Haemostasis*. 1993;**70**(6):1024-1029
- [12] Zwaal RF, Bevers EM. Platelet phospholipid asymmetry and its significance in hemostasis. *Sub-Cellular Biochemistry*. 1983;**9**:299-334
- [13] Alberio L, Safa O, Clemetson KJ, Esmon CT, Dale GL. Surface expression and functional characterization of alpha-granule factor V in human platelets: Effects of ionophore A23187, thrombin, collagen, and convulxin. *Blood*. 2000;**95**(5):1694-1702
- [14] Kulkarni S, Jackson SP. Platelet factor XIII and calpain negatively regulate integrin alphaIIb beta3 adhesive function and thrombus growth. *The Journal of Biological Chemistry*. 2004;**279**(29):30697-30706
- [15] Dale GL, Friese P, Batar P, Hamilton SF, Reed GL, Jackson KW,

- et al. Stimulated platelets use serotonin to enhance their retention of procoagulant proteins on the cell surface. *Nature*. 2002;**415**(6868):175-179
- [16] Remenyi G, Szasz R, Friese P, Dale GL. Role of mitochondrial permeability transition pore in coated-platelet formation. *Arteriosclerosis, Thrombosis, and Vascular Biology*. 2005;**25**(2):467-471
- [17] Panteleev MA, Ananyeva NM, Greco NJ, Ataullakhanov FI, Saenko EL. Two subpopulations of thrombin-activated platelets differ in their binding of the components of the intrinsic factor X-activating complex. *Journal of Thrombosis and Haemostasis*. 2005;**3**(11):2545-2553
- [18] Jobe SM, Wilson KM, Leo L, Raimondi A, Molkentin JD, Lentz SR, et al. Critical role for the mitochondrial permeability transition pore and cyclophilin D in platelet activation and thrombosis. *Blood*. 2008;**111**(3):1257-1265
- [19] Agbani EO, van den Bosch MTJ, Brown E, Williams CM, Mattheij NJA, Cosemans JMEM, et al. Coordinated membrane ballooning and procoagulant spreading in human platelets. *Circulation*. 2015;**132**(15):1414-1424
- [20] Mazepa M, Hoffman M, Monroe D. Superactivated platelets: Thrombus regulators, thrombin generators, and potential clinical targets. *Arteriosclerosis, Thrombosis, and Vascular Biology*. 2013;**33**(8):1747-1752
- [21] Jobe SM, Leo L, Eastvold JS, Dickneite G, Ratliff TL, Lentz SR, et al. Role of FcR γ and factor XIIIa in coated platelet formation. *Blood*. 2005;**106**(13):4146-4151
- [22] Choo H-J, Kholmukhamedov A, Zhou C, Jobe S. Inner mitochondrial membrane disruption links apoptotic and agonist-initiated phosphatidylserine externalization in platelets. *Arteriosclerosis, Thrombosis, and Vascular Biology*. 2017;**37**(8):1503-1512
- [23] Kim J-S, He L, Lemasters JJ. Mitochondrial permeability transition: A common pathway to necrosis and apoptosis. *Biochemical and Biophysical Research Communications*. 2003;**304**(3):463-470
- [24] Hunter DR, Haworth RA, Southard JH. Relationship between configuration, function, and permeability in calcium-treated mitochondria. *The Journal of Biological Chemistry*. 1976;**251**(16):5069-5077
- [25] Bernardi P, Forte M. The mitochondrial permeability transition pore. *Novartis Foundation Symposium*. 2007;**287**:157-164. Discussion 164-169
- [26] Hathaway DR, Adelstein RS. Human platelet myosin light chain kinase requires the calcium-binding protein calmodulin for activity. *Proceedings of the National Academy of Sciences of the United States of America*. 1979;**76**(4):1653-1657
- [27] Shattil SJ, Brass LF. Induction of the fibrinogen receptor on human platelets by intracellular mediators. *The Journal of Biological Chemistry*. 1987;**262**(3):992-1000
- [28] Podoplelova NA, Sveshnikova AN, Kotova YN, Eckly A, Receveur N, Nechipurenko DY, et al. Coagulation factors bound to procoagulant platelets concentrate in cap structures to promote clotting. *Blood*. 2016;**128**(13):1745-1755
- [29] Kholmukhamedov A, Jobe S. Necrotic but not apoptotic platelets are functionally procoagulant. *Blood*. 2018;**132**(Suppl 1):2420
- [30] McArthur K, Chappaz S, Kile BT. Apoptosis in megakaryocytes and

platelets: The life and death of a lineage. *Blood*. 2018;**131**(6):605-610

[31] Pleines I, Lebois M, Gangatirkar P, Au AE, Lane RM, Henley KJ, et al. Intrinsic apoptosis circumvents the functional decline of circulating platelets but does not cause the storage lesion. *Blood*. 2018;**132**(2):197-209

[32] Mason KD, Carpinelli MR, Fletcher JI, Collinge JE, Hilton AA, Ellis S, et al. Programmed anuclear cell death delimits platelet life span. *Cell*. 2007;**128**(6):1173-1186

[33] Josefsson EC, James C, Henley KJ, Debrincat MA, Rogers KL, Dowling MR, et al. Megakaryocytes possess a functional intrinsic apoptosis pathway that must be restrained to survive and produce platelets. *The Journal of Experimental Medicine*. 2011;**208**(10):2017-2031

[34] Prodan CI, Vincent AS, Dale GL. Coated-platelet levels increase with number of injuries in patients with mild traumatic brain injury. *Journal of Neurotrauma*. 2016;**33**(9):818-824

[35] Prodan CI, Stoner JA, Dale GL. Acute hemorrhagic complications are associated with lower coated-platelet levels in non-lacunar brain infarction. *Journal of Thrombosis and Haemostasis*. 2015;**13**(12):2233-2239

[36] Prodan CI, Stoner JA, Dale GL. Lower coated-platelet levels are associated with increased mortality after spontaneous intracerebral hemorrhage. *Stroke*. 2015;**46**(7):1819-1825

[37] Prodan CI, Vincent AS, Dale GL. Coated-platelet levels are persistently elevated in patients with mild traumatic brain injury. *The Journal of Head Trauma Rehabilitation*. 2014;**29**(6):522-526

[38] Kirkpatrick AC, Vincent AS, Dale GL, Prodan CI. Coated-platelets

predict stroke at 30 days following TIA. *Neurology*. 2017;**89**(2):125-128

[39] Kirkpatrick AC, Stoner JA, Dale GL, Rabadi M, Prodan CI. Higher coated-platelet levels in acute stroke are associated with lower cognitive scores at three months post infarction. *Journal of Stroke and Cerebrovascular Diseases*. 2019;**28**(9):2398-2406

[40] Kirkpatrick AC, Tafur AJ, Vincent AS, Dale GL, Prodan CI. Coated-platelets improve prediction of stroke and transient ischemic attack in asymptomatic internal carotid artery stenosis. *Stroke*. 2014;**45**(10):2995-3001

[41] Ray B, Pandav VM, Mathews EA, Thompson DM, Ford L, Yearout LK, et al. Coated-platelet trends predict short-term clinical outcome after subarachnoid hemorrhage. *Translational Stroke Research*. 2018;**9**(5):459-470

[42] Sakurai Y, Hardy ET, Ahn B, Tran R, Fay ME, Ciciliano JC, et al. A microengineered vascularized bleeding model that integrates the principal components of hemostasis. *Nature Communications*. 2018;**9**(1):509

[43] Kholmukhamedov A, Jobe S. Mitochondrial uncoupling prevents procoagulant platelet formation resulting in decreased thrombus stability. *Blood*. 2018;**132**(Suppl 1):2421

[44] Panteleev MA, Saenko EL, Ananyeva NM, Ataullakhanov FI. Kinetics of factor X activation by the membrane-bound complex of factor IXa and factor VIIa. *The Biochemical Journal*. 2004;**381**(Pt 3):779-794

[45] Ataullakhanov FI, Panteleev MA. Mathematical modeling and computer simulation in blood coagulation. *Pathophysiology of Haemostasis and Thrombosis*. 2005;**34**(2-3):60-70

[46] Panteleev MA, Balandina AN, Lipets EN, Ovanesov MV,

- Ataullakhanov FI. Task-oriented modular decomposition of biological networks: Trigger mechanism in blood coagulation. *Biophysical Journal*. 2010;**98**(9):1751-1761
- [47] Ovanesov MV, Ananyeva NM, Pantelev MA, Ataullakhanov FI, Saenko EL. Initiation and propagation of coagulation from tissue factor-bearing cell monolayers to plasma: Initiator cells do not regulate spatial growth rate. *Journal of Thrombosis and Haemostasis*. 2005;**3**(2):321-331
- [48] Sinauridze EI, Kireev DA, Popenko NY, Pichugin AV, Pantelev MA, Krymskaya OV, et al. Platelet microparticle membranes have 50- to 100-fold higher specific procoagulant activity than activated platelets. *Thrombosis and Haemostasis*. 2007;**97**(3):425-434
- [49] Krishnaswamy S, Nesheim ME, Prydzial EL, Mann KG. Assembly of prothrombinase complex. *Methods in Enzymology*. 1993;**222**:260-280
- [50] Mann KG, Krishnaswamy S, Lawson JH. Surface-dependent hemostasis. *Seminars in Hematology*. 1992;**29**(3):213-226
- [51] Gilbert GE, Novakovic VA, Shi J, Rasmussen J, Pipe SW. Platelet binding sites for factor VIII in relation to fibrin and phosphatidylserine. *Blood*. 2015;**126**(10):1237-1244
- [52] Gilbert GE, Arena AA. Activation of the factor VIIIa-factor IXa enzyme complex of blood coagulation by membranes containing phosphatidyl-L-serine. *The Journal of Biological Chemistry*. 1996;**271**(19):11120-11125
- [53] Nesheim ME, Nichols WL, Cole TL, Houston JG, Schenk RB, Mann KG, et al. Isolation and study of an acquired inhibitor of human coagulation factor V. *The Journal of Clinical Investigation*. 1986;**77**(2):405-415
- [54] Maroney SA, Haberichter SL, Friese P, Collins ML, Ferrel JP, Dale GL, et al. Active tissue factor pathway inhibitor is expressed on the surface of coated platelets. *Blood*. 2007;**109**(5):1931-1937
- [55] Nechipurenko DY, Receveur N, Yakimenko AO, Shepelyuk TO, Yakusheva AA, Kerimov RR, et al. Clot contraction drives the translocation of procoagulant platelets to thrombus surface. *Arteriosclerosis, Thrombosis, and Vascular Biology*. 2019;**39**(1):37-47
- [56] Mattheij NJA, Gilio K, van Kruchten R, Jobe SM, Wieschhaus AJ, Chishti AH, et al. Dual mechanism of integrin α IIb β 3 closure in procoagulant platelets. *The Journal of Biological Chemistry*. 2013;**288**(19):13325-13336
- [57] Liu F, Gamez G, Myers DR, Clemmons W, Lam WA, Jobe SM. Mitochondrially mediated integrin α IIb β 3 protein inactivation limits thrombus growth. *The Journal of Biological Chemistry*. 2013;**288**(42):30672-30681
- [58] Yakimenko AO, Verholomova FY, Kotova YN, Ataullakhanov FI, Pantelev MA. Identification of different proaggregatory abilities of activated platelet subpopulations. *Biophysical Journal*. 2012;**102**(10):2261-2269
- [59] Alberio L, Ravanat C, Hechler B, Mangin PH, Lanza F, Gachet C. Delayed-onset of procoagulant signalling revealed by kinetic analysis of COAT platelet formation. *Thrombosis and Haemostasis*. 2017;**117**(6):1101-1114
- [60] Kholmukhamedov A, Jobe S. Mitochondria and platelet cell death. *Thrombosis and Haemostasis*. 2017;**117**(11):2207-2208
- [61] Kholmukhamedov A, Janecke R, Choo H-J, Jobe SM. The mitochondrial

calcium uniporter regulates
procoagulant platelet formation. *Journal
of Thrombosis and Haemostasis*.
2018;**16**(11):2315-2321

[62] Kholmukhamedov A, Jobe S.
Procoagulant platelets get squeezed to
define the boundaries of the hemostatic
plug. *Arteriosclerosis, Thrombosis, and
Vascular Biology*. 2019;**39**(1):5-6